

Editorial Comment

THE TRAUMA HEMOSTASIS AND OXYGENATION RESEARCH NETWORK'S REMOTE DAMAGE CONTROL RESUSCITATION SYMPOSIUM

Philip C. Spinella* and Geir Strandenes†

*Department of Pediatrics, Division of Critical Care, Washington University in St Louis, St Louis, Missouri; and

†Norwegian Naval Special Operation Commando, Bergen, Norway

Received 22 Aug 2013; first review completed 11 Sep 2013; accepted in final form 13 Sep 2013

ABSTRACT—This year, the Trauma Hemostasis and Oxygenation Research Network had its third annual conference from June 17 to 19 at the Solstrand Hotel, near Bergen, Norway. It was sponsored and organized by the Norwegian Naval Special Operation Commando together with the Norwegian Air Ambulance Foundation. The Trauma Hemostasis and Oxygenation Research Network is composed of more than 150 members from 16 countries who all have a common interest in the prevention and treatment of traumatic hemorrhagic shock. The network is multidisciplinary to include members from both the military and civilian medical community representing areas of surgery, critical care, emergency medicine, transfusion medicine, anesthesiology, hematology, and basic science.

KEYWORDS—Trauma, transfusion, prehospital, resuscitation

The mission of the Trauma Hemostasis and Oxygenation Research (THOR) Network is to reduce morbidity and mortality for patients with traumatic hemorrhagic shock in the prehospital phase. Our main focus is to improve resuscitation strategies through research, education, and training (www.rdc.org). The annual Remote Damage Control Resuscitation (RDCR) symposium is the main venue for the network to achieve this goal (1, 2).

The goal of the 2013 RDCR symposium was to focus on training methods to improve implementation of RDCR principles, to present current research on RDCR, and provide practical information from international leaders in trauma resuscitation on their approach to RDCR. During the 3-day symposium, the development of research protocols were discussed and refined. For example, Dr Alan Murdock presented his initial concept of a whole blood compared with blood component pilot trial for patients with massive traumatic bleeding (3). Dr Jean Louis Vincent focused on alternatives to randomized controlled trials to examine the effect of RDCR principles (4), and Dr Mitchell Cohen presented how a systems biology approach can be used to improve outcomes in this population (5). The symposium also allowed for the dissemination of results from previously performed studies to educate our members on the latest science in the field. Dr Andre Cap, director of the Blood Research Division at the US Army Institute of Surgical Research, presented *in vitro* and *in vivo* data that compared platelet function for platelets stored at 4°C versus 22°C (6, 7). There were presentations and roundtable discussions led by international experts in transfusion medicine that highlighted the ongoing paradigm shift from using crystalloids and colloids for hemorrhagic shock to the use of blood products and hemostatic agents in the prehospital arena (8–11). Of particular interest were presentations that included discussions regarding how alternative processing of blood products, for example, by lyophilization of plasma or solvent detergent treatment of large pools of plasma, may improve

availability and safety of plasma products (9,12). Practical information regarding the application of RDCR principles in both military and civilian scenarios as well as for rapid versus prolonged evacuation times was presented by internationally respected subject matter experts from Great Britain, United States, Israel, Austria, Germany, and Norway (8, 10, 13–18). The annual RDCR symposium also provides an opportunity for the first responder whether that is a military medic or a civilian paramedic to describe limitations in the field to apply RDCR principles and to share what new research, products, or devices are necessary for them to improve care in the field. Methods in training prehospital providers were presented during the conference providing evidence in a live demonstration showing the feasibility of collecting and administering fresh whole blood in a far forward scenario (19).

Damage control resuscitation (DCR) concepts have been reported in detail previously, but can be summarized by combining the appropriate use of hypotensive resuscitation and hemostatic resuscitation with the goal of reducing death from shock and coagulopathy (20, 21). Hemostatic resuscitation is the use of a balanced amount of red blood cells, plasma, and platelets to simultaneously and adequately treat or prevent the progression of shock and coagulopathy. As described previously in the supplement for the 2012 RDCR symposium, exporting DCR concepts to the prehospital arena is the essence of RDCR (22). Remote DCR can involve either short or long transport times, which can affect the application of RDCR principles. For example, the application of hypotensive resuscitation, which allows for a limited use of fluids to maintain systolic blood pressure between 80 and 90 mmHg to prevent overresuscitation and continued bleeding, may not be appropriate if applied to patients in austere environments who required prolonged evacuations. During this year's symposium, Dr Kevin Ward led a pro/con debate on this topic.

The ideal prehospital resuscitation strategy for patients with life-threatening bleeding is controversial. There are some data

to guide RDCR principles. Civilian and military trauma reports indicate that the majority of deaths occur before reaching a surgical facility (23–26). Hemorrhage is a major mechanism of death in combat injuries, and death from hemorrhage represents 80% to 90% of those potentially salvageable (27). In the civilian community, trauma represents the leading cause of death in the age from 1 to 44 years, and 50% of deaths within the first 24 h are due to hemorrhage (28). Trauma-induced coagulopathy occurs quickly after injury and is highly lethal (29–31). These facts lead us to the conclusion that one of the most important “gaps” to be filled in the management of the exsanguinating patient appears in the prehospital phase. Making blood products available as far forward as possible might improve survival, especially from noncompressible injuries. In addition, improvements in identifying and monitoring for shock and coagulopathy during the prehospital phase of resuscitation would also likely dramatically impact outcomes in this population.

Many believe the next large breakthrough in improving outcomes for patients with traumatic injuries will come from improvement in prehospital survival (32). The THOR network through the annual Solstrand RDCR symposium is committed to promoting the research, training, and education needed to determine optimal prehospital resuscitation strategies with the ultimate goal of reducing morbidity and mortality from traumatic hemorrhage.

The following supplement is a collection of manuscripts from speakers at the 2013 RDCR symposium. As cochairs of THOR and the RDCR symposium, we are very proud of the quality of manuscripts provided for this supplement by many leaders in the world on this topic of RDCR. We hope that this supplement is informative and stimulates investigation, training, and education in this important clinical topic.

REFERENCES

- Spinella PC, Strandenes G, Rein EB, Seghatchian J, Hervig T: Symposium on fresh whole blood for severe hemorrhagic shock: from in-hospital to far forward resuscitations. *Transfus Apherisis Sci* 46(1):113–117, 2012.
- Strandenes G, Spinella PC: The Solstrand remote damage control resuscitation symposium. *Transfusion* 53(Suppl 1):6S–8S, 2013.
- Murdock AD, Berseus O, Hervig T, Strandenes G, Lunde TH: Whole blood: the future of traumatic hemorrhagic shock resuscitation. *Shock* 41(S1):62–69, 2014.
- Vincent JL, Sakr Y, Lelubre C: The future of observational research and RCTs in transfusion medicine. *Shock* 41(S1):98–101, 2014.
- Jenkins DH, Rappold JF, Badloe JF, Berseus O, Blackburne L, Brohi KH, Butler FK, Cap AP, Cohen MJ, Davenport R, et al.: Trauma hemostasis and oxygenation research position paper on remote damage control resuscitation: definitions, current practice, and knowledge gaps. *Shock* 41(S1):3–12, 2014.
- Pidcoke HF, Spinella PC, Ramasubramanian AK, Strandenes G, Hervig T, Ness PM, Cap AP: Refrigerated platelets for the treatment of acute bleeding: a review of the literature and reexamination of current standards. *Shock* 41(S1):51–53, 2014.
- Reddoch KM, Pidcoke HF, Montgomery RK, Fedyk C, Aden JK, Ramasubramanian AN, Cap AP: Hemostatic function of apheresis platelets stored at 4°C and 22°C. *Shock* 41(S1):54–61, 2014.
- Schött U: Prehospital coagulation monitoring of resuscitation with point-of-care devices. *Shock* 41(S1):26–29, 2014.
- Hervig T, Doughty H, Ness P, Badloe JF, Berseus O, Glassberg E, Heier HE: Prehospital use of plasma: the blood bankers’ perspective. *Shock* 41(S1):39–43, 2014.
- Schochl H, Schlimp CJ, Maegele M: Tranexamic acid, fibrinogen concentrate, and prothrombin complex concentrate: data to support prehospital use? *Shock* 41(S1):44–46, 2014.
- Medby C: Is there a place for crystalloids and colloids in remote damage control resuscitation? *Shock* 41(S1):47–50, 2014.
- Pasquier P, Dubost C, Nau A: Implementation and execution of Military Forward Resuscitation Programs. *Shock* 41(S1):102, 2014.
- Hooper TJ, DePasquale M, Strandenes G, Sunde G, Ward KR: Challenges and possibilities in forward resuscitation. *Shock* 41(S1):13–20, 2014.
- Spinella PC, Doctor A: Role of transfused red blood cells for shock and coagulopathy within remote damage control resuscitation. *Shock* 41(S1):30–34, 2014.
- Moore EE, Chin TL, Chapman MC, Gonzalez E, Moore HB, Silliman CC, Hansen KC, Sauaia A, Banerjee A: Plasma first in the field for postinjury hemorrhagic shock. *Shock* 41(S1):35–38, 2014.
- Strandenes G, Berseus O, Cap AP, Hervig T, Reade M, Prat N, Sailliol A, Gonzales R, Simon CD, Ness P, et al.: Low titer group O whole blood in emergency situations. *Shock* 41(S1):70–75, 2014.
- Jenkins D, Stubbs J, Williams S, Berns K, Zielinski M, Strandenes G, Zietlow S: Implementation and execution of civilian remote damage control resuscitation programs. *Shock* 41(S1):84–89, 2014.
- Hooper TJ, Nadler R, Badloe J, Butler FK, Glassberg E: Implementation and execution of military forward resuscitation programs. *Shock* 41(S1):90–97, 2014.
- Strandenes G, DePasquale M, Cap AP, Hervig TA, Kristoffersen EK, Hickey M, Cordova C, Berseus O, Eliassen HS, Fisher L, et al.: Emergency whole-blood use in the field: a simplified protocol for collection and transfusion. *Shock* 41(S1):76–83, 2014.
- Holcomb JB, Jenkins D, Rhee P, Johannigman J, Mahoney P, Mehta S, Cox ED, Gehrke MJ, Beilman GJ, Schreiber M, et al.: Damage control resuscitation: directly addressing the early coagulopathy of trauma. *J Trauma* 62(2):307–310, 2007.
- Spinella PC, Holcomb JB: Resuscitation and transfusion principles for traumatic hemorrhagic shock. *Blood Rev* 23(6):231–240, 2009.
- Gerhardt RT, Strandenes G, Cap AP, Rentas FJ, Glassberg E, Mott J, Dubick MA, Spinella PC, THOR Network, RemTORN Study Groups: Remote damage control resuscitation and the Solstrand conference: defining the need, the language, and a way forward. *Transfusion* 53(Suppl 1):9S–16S, 2013.
- Acosta JA, Yang JC, Winchell RJ, Simons RK, Fortlage DA, Hollingsworth-Fridlund P, Hoyt DB: Lethal injuries and time to death in a level I trauma center. *J Am Coll Surg* 186(5):528–533, 1998.
- Gerhardt RT, De Lorenzo RA, Oliver J, Holcomb JB, Pfaff JA: Out-of-hospital combat casualty care in the current war in Iraq. *Ann Emerg Med* 53(2):169–174, 2009.
- MacLeod JB, Cohn SM, Johnson EW, McKenney MG: Trauma deaths in the first hour: are they all unsalvageable injuries? *Am J Surg* 193(2):195–199, 2007.
- Demetriades D, Murray J, Charalambides K, Alo K, Velmahos G, Rhee P, Chan L: Trauma fatalities: time and location of hospital deaths. *J Am Coll Surg* 198(1):20–26, 2004.
- Eastridge BJ, Hardin M, Cantrell J, Oetjen-Gerdes L, Zubko T, Mallak C, Wade CE, Simmons J, Mace J, Mabry R, Mace J, Mabry R, Bolenbaucher R, Blackburne LH: Died of wounds on the battlefield: causation and implications for improving combat casualty care. *J Trauma* 71(Suppl 1):S4–S8, 2011.
- Nunez TC, Cotton BA: Transfusion therapy in hemorrhagic shock. *Curr Opin Crit Care* 15(6):536–541, 2009.
- Hess JR, Brohi K, Dutton RP, Hauser CJ, Holcomb JB, Kluger Y, Mackway-Jones K, Parr MJ, Rizoli SB, Yukioka T, et al.: The coagulopathy of trauma: a review of mechanisms. *J Trauma* 65(4):748–754, 2008.
- Brohi K, Singh J, Heron M, Coats T: Acute traumatic coagulopathy. *J Trauma* 54(6):1127–1130, 2003.
- Patregnani JT, Borgman MA, Maegele M, Wade CE, Blackburne LH, Spinella PC: Coagulopathy and shock on admission is associated with mortality for children with traumatic injuries at combat support hospitals. *Pediatr Crit Care Med* 13(3):273–277, 2012.
- Gerhardt RT, Berry JA, Blackburne LH: Analysis of life-saving interventions performed by out-of-hospital combat medical personnel. *J Trauma* 71(Suppl 1):S109–S113, 2011.